

AD-A076 597

CALIFORNIA UNIV BERKELEY OPERATIONS RESEARCH CENTER
SOLUTIONS OF SPHERICAL INEQUALITIES.(U)

F/G 12/1

SEP 79 D GALE

N00014-76-C-0314

UNCLASSIFIED ORC-79-10


NL

| OF |
AD
A076597



END
DATE
FILMED
12-79

DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

ORC 79-10
SEPTEMBER 1979

(12)

F


LEVEL

SOLUTIONS OF SPHERICAL INEQUALITIES

by
DAVID GALE

ADA 076597

DDC FILE COPY



DISTRIBUTION STATEMENT A

Approved for public release
Distribution Unlimited

OPERATIONS
RESEARCH
CENTER

DDC
RECEIVED
NOV 13 1979
A

UNIVERSITY OF CALIFORNIA • BERKELEY

79 11 09 026

SOLUTIONS OF SPHERICAL INEQUALITIES

by

David Gale
Department of Industrial Engineering
and Operations Research
University of California, Berkeley

Accession For	
NTIS CRARI	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist.	Avail and/or special
A	

SEPTEMBER 1979

ORC 79-10

This research has been partially supported by the National Science Foundation under Grant SOC78-05196 and the Office of Naval Research under Contract N00014-76-C-0134 with the University of California. Reproduction in whole or in part is permitted for any purpose of the United States Government.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER (14) ORC-79-18	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) (6) SOLUTIONS OF SPHERICAL INEQUALITIES,		5. TYPE OF REPORT & PERIOD COVERED (9) Research Report, 10
7. AUTHOR(s) (10) David/Gale		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Operations Research Center University of California Berkeley, California 94720		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS National Science Foundation 1800 G Street, N.W. Washington, D.C. 20550		12. REPORT DATE (11) Sep 1979
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (12) 10		13. NUMBER OF PAGES 9
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Also supported by the Office of Naval Research under Contract N00014-76-C-0134.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Quadratic Programming Circumscribed Ball		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) (SEE ABSTRACT) next page 270 750 Gm		

ABSTRACT

↓
It is shown that the problem of finding a point in the intersection of a finite number of balls in n -space is equivalent to solving a very simple (one constraint) quadratic program.
↑

SOLUTIONS OF SPHERICAL INEQUALITIES

by

David Gale

There are many examples of natural geometrical problems which can be solved by linear or quadratic programming. For example the problem of finding the largest ball contained in a given convex polytope (the *inscribed ball*) turns out to be a linear program, while the problem of finding the smallest ball containing a convex polytope (the *circumscribed ball*) is equivalent to a quadratic program [1]. In this note we extend the result of [1] by showing that the following problem can be solved by a simple quadratic program:

Problem:

Given a finite set of balls in R^n decide whether their intersection is nonempty and if so find a point in it.

To formulate the problem algebraically let c_i be the center, ρ_i the radius of the i^{th} ball, $i = 1, \dots, n$.

Problem:

Solve the system of inequalities

$$(1) \quad (x - c_i)^2 \leq \rho_i^2, \quad i = 1, \dots, m,$$

or show that no solution exists.

An obvious modification of this problem is to find a vector x and number z such that

[†]In this paper the scalar product of vectors x and y is written xy the scalar product of x with itself is x^2 .

(2) z is a maximum

subject to

$$(3) \quad (x - c_i)^2 - \rho_i^2 + z \leq 0, \quad i = 1, \dots, m.$$

Letting (\bar{x}, \bar{z}) be the solution of (2), (3), it is clear that (1) has a solution if and only if $\bar{z} \geq 0$. For the special case $\rho_i = 0$ for all i it is also clear that \bar{z} is negative (unless all c_i are equal) in which case, by definition, \bar{x} is the center and $-\bar{z}$ is the radius of the circumscribed ball.

The point of this note is to show that the problem (2), (3) has a dual which is a very simple quadratic program and hence can be solved by well known techniques. It then follows that if the c_i and ρ_i are rational then (2), (3) have a rational solution which is perhaps not immediately obvious.

Our result is the following. Consider the quadratic program of finding a nonnegative m -vector $\lambda = (\lambda_1, \dots, \lambda_m)$ and a number μ such that

(2)* μ is a minimum

subject to

$$(3)^* \quad \mu = \left(\sum \lambda_i c_i \right)^2 + \sum \lambda_i (\rho_i^2 - c_i^2)$$

and

$$\sum \lambda_i = 1 \quad \left(\sum \text{ is short for } \sum_{i=1}^m \right).$$

We assert that (2), (3) and $(2)^*$, $(3)^*$ are dual problems. More precisely:

Theorem:

Let $(\bar{\lambda}, \bar{\mu})$ solve $(2)^*$, $(3)^*$. Then $(\bar{x}, \bar{\mu})$ solves (2), (3) where $\bar{x} = \sum \bar{\lambda}_i c_i$.

■ We first prove the usual inequality (so called "weak duality"). Let (x, z) be any solution of (3) and (λ, μ) any solution of $(3)^*$. Multiplying (3) by λ_i and summing gives

$$\begin{aligned} 0 &\leq \sum \lambda_i (x - c_i)^2 - \sum \lambda_i \rho_i^2 + z = x^2 - 2 \left(\sum \lambda_i c_i \right) x + \sum \lambda_i (c_i^2 - \rho_i^2) + z \\ &= \left(x - \sum \lambda_i c_i \right)^2 - \left(\sum \lambda_i c_i \right)^2 - \sum \lambda_i (\rho_i^2 - c_i^2) + z \\ &= \left(x - \sum \lambda_i c_i \right)^2 + z - \mu \geq z - \mu, \text{ (from } (3)^*) \end{aligned}$$

so

$$(4) \quad z \leq \mu.$$

Now suppose $(\bar{\lambda}, \bar{\mu})$ is a solution of $(2)^*$, $(3)^*$, and let $\bar{x} = \sum \bar{\lambda}_i c_i$. The Kuhn-Tucker conditions for this problem assert the existence of a number u such that

$$\begin{aligned} (5) \quad u &\leq 2\bar{x}c_i + \rho_i^2 - c_i^2 \text{ for all } i \text{ and} \\ u &= 2\bar{x}c_i + \rho_i^2 - c_i^2 \text{ if } \bar{\lambda}_i > 0. \end{aligned}$$

Multiplying (5) by $\bar{\lambda}_i$ and summing gives

$$u = 2\bar{x}^2 + \sum \lambda_i (\rho_i^2 - c_i^2),$$

or

$$(6) \quad u - \bar{x}^2 = \bar{x}^2 + \sum \lambda_i (\rho_i^2 - c_i^2) = \bar{\mu} \text{ from (3)}^*.$$

Subtracting \bar{x}^2 from (5) gives

$$u - \bar{x}^2 \leq \rho_i^2 - (\bar{x} - c_i)^2$$

so from (6)

$$(\bar{x} - c_i)^2 - \rho_i^2 + \bar{\mu} \leq 0$$

so $(\bar{x}, \bar{\mu})$ is a solution of (3) and from (4) it is optimal, completing the proof. ■

Remarks:

1. The fact that problems (2), (3) and $(2)^*$, $(3)^*$ are nonlinear duals can be derived by appealing to general nonlinear duality theorems, e.g., [2] and making some further simplifications. For this special case, however, we prefer the self-contained proof which uses only the standard optimality criterion (Kuhn-Tucker conditions) for quadratic programs.

2. One might hope to generalize the result of this note by replacing the spherical inequalities of (1) by general positive definite inequalities so that the spheres of (1) become ellipsoids. The following interesting example due to G. Bergman shows that there is no hope of doing this. The following inequalities describe ellipses in the plane.

$$2x^2 + 3xy + 2y^2 - 4x - 4y + 2 \leq 0$$

$$2x^2 + 2xy + y^2 - 2x \leq 0$$

$$x^2 + xy + y^2 - y \leq 0.$$

These inequalities have the unique solution $(2^{2/3} - 2^{1/3}, 2 - 2^{2/3})$.

3. The above example involves three inequalities. It seems possible that for the case of only two positive definite inequalities rational solutions will always exist. The answer is not known at this time.

BIBLIOGRAPHY

- [1] Elzinga, D. J. and D. W. Hearn, "The Minimum Covering Sphere Problem," Management Science, Vol. 19, No. 1, pp. 96-140, (September 1972).
- [2] Wolfe, P., "A Duality Theorem for Nonlinear Programming," Quarterly of Applied Mathematics, Vol. 19, pp. 239-244, (1961).